

## Allometric models for non-destructive leaf area estimation in *Eugenia uniflora* (L.)

### Modelos alométricos para la estimación no destructiva del área foliar en *Eugenia uniflora* (L.)

Pompelli, M. F.<sup>1\*</sup>, Figueirôa, J. M.<sup>2</sup>, Lozano-Isla, F.<sup>1</sup>.

\* Corresponding author. [marcelo.pompelli@ufpe.br](mailto:marcelo.pompelli@ufpe.br)

#### Abstract

We aimed to propose a reliable and accurate model using non-destructive measurements of leaf length (L) and/or width (W) for estimating leaf area (LA) of Surinam cherry (*Eugenia uniflora* L.). For model construction, 560 leaves were randomly sampled from different levels of the tree canopies and encompassed the full spectrum of measurable leaf sizes. Power models better fit *E. uniflora* leaf area than linear models; but, among of them, the best fit were made when product of the L and W (LW) were used. To validate these models, independent data set of 156 leaves were used. Thus, we developed a single power model ( $Y_i = \beta_0 x^{\beta_1}$ ) [LA = 0.685 (LW)<sup>0.989</sup>; standard errors:  $\beta_0 = 0.014$ ,  $\beta_1 = 0.005$ ;  $R_a^2 = 0.997$ ] with high precision and accuracy, random dispersal pattern of residuals and unbiased. A simpler linear model [LA = 0.094 + (LW \* 0.655); standard errors:  $\beta_0 = 0.025$ ,  $\beta_1 = 0.001$ ;  $R_a^2 = 0.998$ ] also described here to estimate leaf area of *E. uniflora*, which are as good as the first. The simplicity of the latter model may be relevant in field studies, as it does not demand high precision or expensive instruments.

**Keywords:** Surinam cherry; estimate model; leaf length; leaf width

#### Resumen

Nuestro objetivo fue proponer un modelo confiable y preciso utilizando mediciones no destructivas de la longitud de la hoja (L) y / o el ancho (W) para estimar el área foliar (LA) de la cereza de Surinam (*Eugenia uniflora* L.). Para la construcción del modelo, se tomaron 560 hojas al azar de diferentes niveles de las copas de los árboles y abarcaron todo el espectro de tamaños de hojas medibles. Los modelos exponenciales se ajustan mejor al área foliar de *E. uniflora* que los modelos lineales; el mejor ajuste se realizó cuando se usaron productos de L y W (LW). Para validar estos modelos, se utilizaron conjuntos de datos independientes de 156 hojas. Por lo tanto, desarrollamos un único modelo de potencia ( $Y_i = \beta_0 x^{\beta_1}$ ) [LA = 0.685 (LW)<sup>0.989</sup>; error estándar:  $\beta_0 = 0.014$ ,  $\beta_1 = 0.005$ ;  $R_a^2 = 0.997$ ] con alta precisión y exactitud, patrón de dispersión aleatorio de residuales e imparcial. Un modelo lineal más simple [LA = 0.094 + (LW \* 0.655); error estándar:  $\beta_0 = 0.025$ ,  $\beta_1 = 0.001$ ;  $R_a^2 = 0.998$ ] también se describe aquí para estimar el área foliar de *E. uniflora*, que es tan buena como la primera. La simplicidad de este último modelo puede ser relevante en los estudios de campo, ya que no exige instrumentos costosos o de alta precisión.

**Palabras llave:** cereza de Surinam; modelo de estimación; longitud de la hoja; ancho de la hoja

#### Introduction

Leaf area (LA) is one of the six most important traits that drive plant form and function (Díaz *et al.*, 2016). This descriptor has been widely used to describe a range of variables including growth, productivity, photosynthetic efficiency, soil characteristics including salinity and acidity, transfer and exchange of heat, carbon, nutrients and water, which in turn affect plant yield (Cristofori *et al.*, 2007; Pompelli *et al.*, 2012). Thus, a correct determination of the LA becomes even more important in crop species, since the leaf is the organ of greater influence with the environment and it is through this that the agronomic

studies are based on the important decision making.

Directly measure of LA of individuals is both, laborious, expensive as well as a time consuming task and often constrained by logistical factors. Leaf area is traditionally quantified by direct methods, which are destructively or obtained through high-cost equipment, such as AM350 portable leaf area meter (ADC BioScientific Ltd., Hoddesdon, UK). With the intensification of modeling techniques, numerous studies have proposed allometric models to predict the LA of different species (Blanco and Folegatti, 2005; Antunes *et al.*, 2008; Pompelli *et al.*, 2012; Keramatlou *et al.*, 2015; Liu *et al.*, 2017). Hence,

<sup>1</sup>Plant Ecophysiology Laboratory, Federal University of Pernambuco, Department of Botany, CCB, Recife, PE, Brazil, 50670901

<sup>2</sup>Secretaria de Meio Ambiente do Estado de Pernambuco, Recife, Pernambuco, Brazil

simple linear measurements like leaf length (L), leaf width (W) are used in allometric equations to predict the leaf area (Peksen, 2007). Given the fact that *Eugenia uniflora* is a commercial crop species, non-destructive methods such as allometry are best suited for measure leaf area because preserve the leaf on the plant instead destructive methods (Cristofori *et al.*, 2007). This paper describes, for first time allometric equations to predict leaf area of *Eugenia uniflora* checking the accuracy and perform an unbiased tool for use in land and agronomic studies.

## Material and Methods

For model construction, 560 healthy leaves were collected at least 20 healthy plants naturally grown at Caetés Ecological Station, Paulista, Pernambuco, Brazil (7°55'28"S; 34°56'02"O; 88 m.a.s.l.) in April 2018 (end of the growing season). To validate the model, an independent data set of 156 leaves were sampled randomly from different levels of the tree canopy, removed from the branches and taken to the laboratory. The maximum leaf length (L) (from lamina tip to the point of the petiole intersection to the midrib) and leaf width (W) (the widest linear length perpendicular to the midrib) were measured to the nearest of 0.001 cm. The leaves were scanned using a scanner (Epson 1200 x 1200 dpi) and images were analyzed using the Image-Pro® Plus software (2001). The leaves encompassed the broadest range as possible. The minimum leaf area sampled was 0.17 cm<sup>2</sup> and maximum was 72.40 cm<sup>2</sup> (Table 1).

Nine theoretical models (more widely used in the literature) were tested, based on different combinations between the components of LA (dependent variable) and respective values of L and W (independent variables). The equations were deduced by the principle of parsimony

(Steel and Penny, 2000), and thus from the “simplest”, or an “optimal” description of the data. All equations were adjusted following the linear simple, modified linear (from exclude  $\beta_0$ ) and power models (more information, see Table 2). All parameters of each model were obtained using DataFit version 8.0.32 (Oakdale Engineering, 2002). The statistical criteria used to select the models were based on (i) the analysis of variance (F test,  $P < 0.001$ ), (ii) adjusted coefficient of determination ( $R_a^2$ ), (iii) mean squared error (MSE), (iv) Student’s t-test ( $P < 0.001$ ) for absolute mean of errors with confidence intervals (Cumming *et al.*, 2007), (v) dispersion pattern of residuals in percentage terms (%) and the best relationship (major  $R_a^2$ ) between observed leaf area and estimated leaf area of the independent data set used to validating equations. The dispersion of the residues were observed in the total sample set, both in small leaves and in larger leaves. The hypothesis of normality of the errors were evaluated, so that heteroscedasticity was considered a reason for model disqualification. These procedures allowed us to assess the occurrence of bias and accuracy in all proposed models (Walther and Moore, 2005).

## Results and Discussion

All nine developed equations (Table 3) presented good predictors of the *E. uniflora* leaf area, since  $R_a^2$  was always higher than 0.8; coefficient higher than those proposed to some crop plants (Cristofori *et al.*, 2007; Peksen, 2007; Kumar, 2009; Souza and Amaral, 2015) and within the range of those proposed by others (Blanco and Folegatti, 2005; Antunes *et al.*, 2008; Demirsoy, 2009; Pompelli *et al.*, 2012; Shabani and Sepaskhah, 2017). Thus, at first glance, all proposed equations should be able to predict with accuracy the leaf area of *E. uniflora*. However, when we analyzed the deviation between estimated leaf area

**Table 1.** Means  $\pm$  standard deviations (SD), minimum (min), maximum (max) values for the leaf length (L), width (W), and leaf area (LA) of the *Eugenia uniflora* L.

L (cm)			W (cm)			LA (cm <sup>2</sup> )		
Mean $\pm$ SD	Min	Max	Mean $\pm$ D	Min	Max	Mean $\pm$ SD	Min	Max
3.55 $\pm$ 2.32	0.72	14.31	1.97 $\pm$ 1.42	0.27	7.62	5.83 $\pm$ 10.73	0.17	72.40

**Table 2.** Statistical models and equations to predict leaf area as a function of linear dimensions of leaves

(Model #1)	Linear	$Y_i = \beta_0 + \beta_1 * \text{Length} + \epsilon_i$
(Model #2)	Linear	$Y_i = \beta_0 + \beta_1 * \text{Width} + \epsilon_i$
(Model #3)	Linear	$Y_i = \beta_0 + \beta_1 * (\text{Length} * \text{Width}) + \epsilon_i$
(Model #4)	Linear without intercept	$Y_i = \beta_1 * \text{Length} + \epsilon_i$
(Model #5)	Linear without intercept	$Y_i = \beta_1 * \text{Width} + \epsilon_i$
(Model #6)	Linear without intercept	$Y_i = \beta_1 * (\text{Length} * \text{Width}) + \epsilon_i$
(Model #7)	Power	$Y_i = \beta_0 * \text{Length}^{\beta_1} + \epsilon_i$
(Model #8)	Power	$Y_i = \beta_0 * \text{Width}^{\beta_1} + \epsilon_i$
(Model #9)	Power	$Y_i = \beta_0 * (\text{Length} * \text{Width})^{\beta_1} + \epsilon_i$

$Y_i$  = leaf area,  $\beta_0$  and  $\beta_1$  = model coefficients and  $\epsilon_i$  = random error

and observed leaf area (Fig. 1), we demonstrate that equations #4, #5, and #8 were biased, because a significant underestimation of LA. In this case, the underestimation can lead for 53.1% of the estimated leaf area (using equation #8) to 93.2% of the estimated leaf area (using equation #4). In similar manner, the equation #6 can lead to overestimation leaf area in about 19.1% of the leaves. All these equations had an estimated significant difference from zero (biased) and were excluded


from the further analysis as recommended by Antunes *et al.* (2008) for coffee trees, by Pompelli *et al.* (2012) for purging-nut trees and by Yadav *et al.* (2007); all agronomic species.

With the disqualification of equations #4, #5, #6 and #8, five equations remained to be analyzed in more details. Thus, a deep analysis of relationship between estimated leaf area and dispersal pattern of residuals, revealed that equations #1 lead to an overestimation of 34.5% and #2 to an underestimation of 36.2% of the leaves (when considering relative errors  $\geq 40\%$ ). A possible source of error must be due to negative values of  $\beta_0$ , that in these equations were bigger than -7.4. Higher values of  $\beta_0$  were previously (Blanco and Folegatti, 2005; Cristofori *et al.*, 2007; Antunes *et al.*, 2008; Pompelli *et al.*, 2012; Schmildt *et al.*, 2015) used to disqualification the allometric equations, because such equations return negative values of LA (*i.e.*, invalid biological condition).

The heteroscedasticity of residuals of model #7, was used to disqualify it, because this model lead to overestimation or underestimation on 21.2% of the leaves (when considering relative errors  $\geq 20\%$ ). The biased pattern of residuals showed in equations #1, #2 and #7 are as large as the smaller sampled leaves. So, we can only recommended the use of these equations when a stratification of the leaf size classes is performed, simultaneously checking the dispersion pattern of the residues in all classes of leaves, as suggested by others (Walther and Moore, 2005; Antunes *et al.*, 2008; Zuur *et al.*, 2010; Pompelli *et al.*, 2012). However, this “solution”, sometimes becomes laborious and impractical, despite the ease of adjustment and operation of this type of model.

From nine initial proposed models, only two models (#3 and #9) were entirely approved. As shows in Fig. 2, the equations #3 lead to overestimation some leaf areas. Therefore, this overestimation is lower than 4%, which

cannot invalidate this equation. In this issue, we argue that the best equation is equation #9, made with power model. Even if the model #7 has been previously disqualified, we can verify that it presents an excellent fit curve ( $R^2_a = 0.985$ ) between linear dimensions of leaves and observed leaf area (Fig. 3B). Models using single leaf dimension power model incorporating L or W may be an interesting option because



**Figure 1.** Statistical analysis of the deviation of the estimated area from the observed area for an individual leaf. Leaf area for *Eugenia uniflora* was estimated using several models in which  $\beta_0$  and  $\beta_1$  are coefficients. Vertical bars denote means and spreads denote 99% confidence intervals of the difference. Numbers below the graph denote model numbers (see further details in the Table 1). Asterisks in the bars denote a biased model.

it requires measurement of only one leaf dimension, thus simplifying measurement procedures (Blanco and Folegatti, 2005; Cristofori *et al.*, 2007; Antunes *et al.*, 2008; Pompelli *et al.*, 2012). Because of this, we prefer to keep this equation to next step of validation.


When we pooled the relationship between observed leaf

area and linear estimated leaf area of the independent sample leaves (Fig. 3, right panel), we verified that equations #3, #7, and #9 returns good fit curves, showing the coefficient of determination above of 0.960. However, as suggested above, the equation #7 returns the lesser determination coefficient of all, besides being present the lesser *P* value ( $P = 0.186$ ) than others ( $P \geq 0.674$ ).

Finally, we argue that from nine initial proposed models, only two models (Model #3 and #9) can provide an unbiased estimation of leaf area using the linear dimensions of leaves. These models were approved in all statistical analysis and then are able to use without errors, both in field and in greenhouses evaluations. However, among then, the

**Table 3.** Statistical models, regression coefficients ( $\beta_0$ ,  $\beta_1$ ), degrees of freedom of residuals (R-d.f.), mean squared error (MSE), coefficients of determination adjusted for the degrees of freedom ( $R^2_{adj}$ ) and *P* value as a function of linear dimensions of leaves.

Models	Coefficients		R-d.f.	MSE	$R^2_{adj}$	P
	$\beta_0$	$\beta_1$				
#1	-7.641	3.854	352	9.311	0.884	<0.001
#2	-7.418	6.834	352	10.207	0.873	<0.001
#3	0.094	0.655	352	0.141	0.998	<0.001
#4	-	2.353	353	23.516	0.819	<0.001
#5	-	4.221	353	23.770	0.818	<0.001
#6	-	0.658	353	0.146	0.996	<0.001
#7	0.407	1.944	352	1.166	0.985	<0.001
#8	2.343	1.591	352	6.198	0.923	<0.001
#9	0.685	0.989	352	0.136	0.997	<0.001




**Figure 2.** The relationship between estimated leaf area and dispersal pattern of residuals to each selected equations. Red oval shape denote strong underestimated (A, B) or overestimated (C) leaf area. Red arrows denote strong biased estimated leaf areas, mainly in the smaller leaves, while black arrows denote slightly skewed estimated leaf areas. See further details in the text.

equation #3 is simpler than equation #9, because it does not require more complex calculations. In the other hand, if the researcher has notion of the error merged in equation #7 and this may lead to an overestimation of approximately 21% of the estimated leaf area, this equation could also be an interesting option because it requires measurement of only one leaf dimension, simplifying measurement procedures (Blanco and Folegatti, 2005; Pompelli *et al.*, 2012), an important aspect specially in the field when a large number of leaves has to be monitored.

## Conclusion

In this study, we developed a reliable and accurate equation to estimate the leaf area or *Eugenia uniflora* using non-destructive method. A power equation, type  $Y_i = \beta_0 x^{\beta_1}$  [LA =  $0.685 (LW)^{0.989}$ ] was made. This equation may estimate



**Figure 3.** The relationship between observed leaf area and linear dimensions of leaf (A, B and C) or between estimated leaf area (D, E and F) for model #3 (A and D), model #7 (B and E) and model #9 (C and F). L, leaf length; LW, product of leaf length and leaf width; ns, not significant;  $R_a^2$ , coefficients of determination adjusted for the degrees of freedom.

the leaf area with 99.7% of accuracy. The simplification of this equation could be done using a linear equation [LA =  $0.094 + (LW * 0.655)$ ] without loss of accuracy. This procedure should be less laborious because use a linear equation instead a power equation. This is the first study that describes with great accuracy an allometric equation to estimate the leaf area of *Eugenia uniflora*, showing all common mistakes in allometric equations published until now.

## Acknowledgements

The authors thank to Secretaria de Meio Ambiente do Estado de Pernambuco (Grants 03/2013) and the

Foundation for Science and Technology of Pernambuco, FACEPE (Grants APQ-0239-2.03/15) for financially supporting this research.

## References

- Antunes, W.C., Pompelli, M.F., Carretero, D.M. and DaMatta, F.M. (2008). Allometric models for non-destructive leaf area estimation in coffee (*Coffea arabica* and *Coffea canephora*). *Annals of Applied Biology*, 153(1), 33-40. <https://doi.org/10.1111/j.1744-7348.2008.00235.x>
- Blanco, F.F. and Folegatti, M.V. (2005). Estimation of leaf area for greenhouse cucumber by linear measurements under salinity and grafting. *Scientia Agricola*, 62(4), 305-309. <http://dx.doi.org/10.1590/S0103-90162005000400001>
- Cristofori, V., Rouphael, Y., Mendoza-de Gyves, E. and Bignami, C. (2007). A simple model for estimating leaf area of hazelnut from linear measurements. *Scientia Horticulturae*, 113(2), 221-225. <https://doi.org/10.1016/j.scienta.2007.02.006>
- Cumming, G., Fidler, F. and Vaux, D.L. (2007). Error bars in experimental biology. *Journal of Cell Biology* 177(1), 7-11. DOI: 10.1083/jcb.200611141
- DataFit version 8.032. (2002). Oakdale Engineering, Oakdale, CA, USA
- Demirsoy, H. (2009). Leaf area estimation in some species of fruit tree by using models as a non-destructive method. *Fruits*, 64(1), 45-51. <https://doi.org/10.1051/fruits/2008049>
- Díaz, S., Kattge, J., Cornelissen, J.H.C., Wright, I.J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C., Colin Prentice, I., Garnier, E., Bönsch, G., Westoby, M., Poorter, H., Reich, P.B., Moles, A.T., Dickie, J., Gillison, A.N., Zanne, A.E., Chave, J., Joseph Wright, S., Sheremet'ev, S.N., Jactel, H., Baraloto, C., Cerabolini, B., Pierce, S., Shipley, B., Kirkup, D., Casanoves, F., Joswig, J.S., Günther, A., Falczuk, V., Rüger, N., Mahecha, M.D. and Gorné, L.D. (2016). The global spectrum of plant form and function. *Nature*, 529, 167-171. <https://doi.org/10.1038/nature16489>
- Image Pro Plus version 4.5.029. (2001). Media Cybernetics Inc. Rockville, MD, USA
- Keramatlou, I., Sharifani, M., Sabouri, H., Alizadeh, M. and Kamkar, B. (2015). A simple linear model for leaf area estimation in Persian walnut (*Juglans regia* L.). *Scientia Horticulturae*, 184, 36-39. <https://doi.org/10.1016/j.scienta.2014.12.017>
- Kumar, R. (2009). Calibration and validation of regression model for non-destructive leaf area estimation of saffron (*Crocus sativus* L.). *Scientia Horticulturae*-Amsterdam 122(1), 142-145. <https://doi.org/10.1016/j.scienta.2009.03.019>
- Liu, Z., Zhu, Y., Li, F. and Jin, G. (2017). Non-destructively predicting leaf area, leaf mass and specific leaf area based on a linear mixed-effect model for broadleaf species. *Ecological Indicators*, 78, 340-350. <https://doi.org/10.1016/j.ecolind.2017.03.025>
- Peksen, E. (2007). Non-destructive leaf area estimation model for faba bean (*Vicia faba* L.). *Scientia Horticulturae*-Amsterdam 113(4), 322-328. <https://doi.org/10.1016/j.scienta.2007.04.003>
- Pompelli, M.F., Antunes, W.C., Ferreira, D.T.R.G., Cavalcante, P.P.G.S., Wanderley-Filho, H.C.L. and Endres, L. (2012). Allometric models for non-destructive leaf area estimation of the *Jatropha curcas*. *Biomass and Bioenergy*, 36, 77-85. <https://doi.org/10.1016/j.biombioe.2011.10.010>
- Schmidt, E.R., Amaral, J.T., Santos, J.S. and Schmidt, O. (2015). Allometric model for estimating leaf area in clonal varieties of coffee (*Coffea canephora*). *Revista Ciência Agronômica*, 46(4), 740-748. <http://dx.doi.org/10.5935/1806-6690.20150061>
- Shabani, A. and Sepaskhah, A.R. (2017). Leaf area estimation by a simple and non-destructive method. *Iran Agricultural Research*, 36(2), 101-104. DOI: [10.22099/IAR.2017.4157](https://doi.org/10.22099/IAR.2017.4157)
- Souza, M.C. and Amaral, C.L. (2015). Non-destructive linear model for leaf area estimation in *Vernonia ferruginea* Less. *Brazilian Journal of Biology*, 75(1), 152-156. <http://dx.doi.org/10.1590/1519-6984.09813>
- Steel, M. and Penny, D. (2000). Parsimony, likelihood, and the role of models in molecular phylogenetics. *Molecular Biology and Evolution*, 17(6), 839-850. DOI: 10.1093/oxfordjournals.molbev.a026364
- Walther, B.A. and Moore, J.L. (2005). The concepts of bias, precision and accuracy, and their use in testing the performance of species richness estimators, with a literature review of estimator performance. *Ecography*, 28(6), 815-829. <https://doi.org/10.1111/j.2005.0906-7590.04112.x>
- Yadav, S.K., Mishra, Y.D. and Singh, R.K. (2007). Total leaf area estimation of *Flemingia semialata* Roxb. by linear regression. *Agricultural Science Digest* 27(1), 44-46.
- Zuur, A.F., Ieno, E.N. and Elphick, C.S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution* 1(1), 3-14. <https://doi.org/10.1111/j.2041-210X.2009.00001.x>